

Some Applications of Spanning Trees in $K_{s,t}$

L.H. Clark, A.T. Mohr, and T.D. Porter*

Department of Mathematics
Southern Illinois University
Carbondale, IL 62901-4408
tporter@math.siu.edu

Abstract

We partition the set of spanning trees contained in the complete graph K_n into spanning trees contained in the complete bipartite graph $K_{s,t}$. This classification shows that some properties of spanning trees in K_n can be derived from trees in $K_{s,t}$. We use Abel's binomial theorem and the formula for spanning trees in $K_{s,t}$ to obtain a proof of Cayley's theorem using partial derivatives. Some results concerning non-isomorphic spanning trees are presented. In particular we count these trees for Q_3 and the Petersen graph.

Keywords. Abel's binomial theorem, Cayley's theorem, hypercube, Petersen graph, spanning trees

1 Introduction

We use the standard notation and terminology which can be found, e.g., in [12]. Let $\tau(G)$ denote the number of labelled spanning trees in a graph G . Let K_n denote the complete graph of n vertices and $K_{s,t}$ the complete bipartite graph with partite sets containing s and t vertices, respectively. It is well known, as in e.g. [2, 3, 4, 5, 6, 10] that

$$\tau(K_n) = n^{n-2}, \quad n \geq 2 \quad (1)$$

$$\tau(K_{s,t}) = s^{t-1}t^{s-1}, \quad s, t \geq 1. \quad (2)$$

We remark that (1) is often referred to as Cayley's theorem. Let $s + t = n$, where $1 \leq s \leq t$. We have the following observation:

Theorem 1. *With $n \geq 2$, any spanning tree T in K_n is a spanning tree in $K_{s,t}$ for a unique pair (s, t) , with $1 \leq s \leq t$ and $s + t = n$.*

*Contact author.

Proof. Consider a spanning tree T in K_n . Because T is a connected bipartite graph it is uniquely 2-colorable. So construct this unique bipartition by properly 2-coloring the vertex set of T with colors red (R) and blue (B). Let the number of red vertices be s and the number of blue vertices be t ; w.l.o.g., let $s \leq t$. We then have T is a spanning tree in this $K_{s,t}$. \square

The converse is straightforward.

Theorem 2. *With $s + t = n$, any spanning tree in $K_{s,t}$ is a spanning tree in K_n .*

Proof. This follows since $K_{s,t}$ is a spanning subgraph of K_n . \square

Theorem 3.

$$\sum_{s=1}^{n-1} \binom{n}{s} \tau(K_{s,n-s}) = 2\tau(K_n).$$

Proof. By combining Theorems 1 and 2 we see that to find $\tau(K_n)$ we can enumerate all labelled spanning trees in the possible $K_{s,t}$ graphs. The double count occurs from the 2-colorings of the partite sets. \square

We now proceed to show the LHS of Theorem 3 implies the RHS yielding a calculus based proof of Cayley's theorem. We shall apply Abel's binomial formula, [1], which states that for any x, y , and z that:

$$(x + y)^n = \sum_{s=0}^n \binom{n}{s} x(x - sz)^{s-1} (y + sz)^{n-s}. \quad (3)$$

Theorem 4. $\tau(K_{s,t}) \implies \tau(K_n)$. *In words, the formula for $\tau(K_n)$ can be derived from the formula for $\tau(K_{s,t})$.*

Proof. From (3) we have

$$n(x + y)^{n-1} = \frac{\partial}{\partial x} (x + y)^n = \sum_{s=0}^n \binom{n}{s} s(x - z)(x - sz)^{s-2} (y + sz)^{n-s} \quad (4)$$

and consequently,

$$\begin{aligned} \frac{\partial^2}{\partial y \partial x} &= n(n-1)(x + y)^{n-2} \\ &= \sum_{s=0}^n \binom{n}{s} s(x - z)(x - sz)^{s-2} (n-s)(y + sz)^{n-s-1}. \end{aligned} \quad (5)$$

We also have,

$$n(x + y)^{n-1} = \frac{\partial}{\partial y} (x + y)^n = \sum_{s=0}^n \binom{n}{s} x(x - sz)^{s-1} (n-s)(y + sz)^{n-s-1}. \quad (6)$$

By substituting $x = n$, $y = 0$, and $z = 1$ into (5) and (6), we obtain, respectively, (7) and (8):

$$n^{n-1} = \sum_{s=1}^n \binom{n}{s} s^{n-s} (n-s)^{s-1} \quad (7)$$

$$n^{n-1} = \sum_{s=1}^n \binom{n}{s} (n-s)^s s^{n-s-1}. \quad (8)$$

Adding (7) to (8) gives

$$2n^{n-1} = \sum_{s=1}^n \binom{n}{s} s^{n-s-1} (n-s)^{s-1} n, \quad (9)$$

which yields the equation in Theorem 3. This gives a proof of Cayley's theorem using partial derivatives. \square

The identity in (9) can also be found in [2, 8, 11]. The ideas in Theorems 1 and 2 are also valid when graphs are unlabelled, since the unique bipartition aspect is a structural property of the graph T . So, for a connected graph G , let $I(G)$ be the number of non-isomorphic spanning trees in G . We have:

Theorem 5. $I(K_n) = \sum_{s=1}^{\lfloor n/2 \rfloor} I(K_{s,n-s}).$ \square

A formula for $I(K_{s,t})$ would then give a formula for $I(K_n)$. We wrote a computer program that generates the set of labelled spanning trees in a graph G . It then partitions this set into its isomorphism classes. Table 1 gives some results found when $G = K_{s,t}$, $I(K_{5,5})$ being the largest calculation in terms of computing time we have been able to produce. The top number in row s and column t corresponds to $\tau(K_{s,t})$ and the bottom number is $I(K_{s,t})$. We have not seen these numbers in Table 1 in the literature.

Observational examples of Theorem 5 and Table 1, using well known values of some $I(K_n)$, are:

$$\begin{aligned} I(K_6) &= 6 = I(K_{1,5}) + I(K_{2,4}) + I(K_{3,3}) \\ &= 1 + 2 + 3, \end{aligned}$$

$$\begin{aligned} I(K_7) &= 11 = I(K_{1,6}) + I(K_{2,3}) + I(K_{3,4}) \\ &= 1 + 3 + 7, \end{aligned}$$

and

$$\begin{aligned} I(K_{10}) &= 106 = I(K_{1,9}) + I(K_{2,8}) + I(K_{3,7}) + I(K_{4,6}) + I(K_{5,5}) \\ &= 1 + 4 + 19 + 45 + 37. \end{aligned}$$

We would like to derive a general or asymptotic formula for $I(K_{s,t})$. An asymptotic formula for $I(K_n)$ is given by Otter [9]

$$I(K_n) \sim pn^{-5/2}r^{-n}, \quad \text{where } p \text{ and } r \text{ are constants.}$$

	1	2	3	4	5	6	7	8	...	n
1	1	1	1	1	1	1	1	1		1
	1	1	1	1	1	1	1	1		1
2	...	4	12	32	80	192	448	1042		
	...	1	2	2	3	3	4	4		
3	81	432	2025	8748	35721	139968		
	3	7	10	14	19	24		
4	4096	32000	221184				
	9	28	45				
5	390625					
	37					

Table 1: Values of τ and I for $K_{s,t}$

Our work so far has given partition numbers for $I(K_{2,n})$ and $I(K_{3,n})$. Let $p_k(n)$ denote the number of partitions of an integer n into k or fewer parts. Then, we have:

$$I(K_{2,n}) = p_2(n-1) = \left\lfloor \frac{n-1}{2} \right\rfloor + 1, \quad \text{for } n \geq 2 \quad (10)$$

$$I(K_{3,n}) = p_3(n-1) + \sum_{k=0}^{n-2} p_2(n-2-k), \quad \text{for } n \geq 4. \quad (11)$$


In (11), we adopt the convention that $p_2(0) = 1$. As an example of (11),

$$\begin{aligned} I(K_{3,5}) &= 10 = p_3(4) + p_2(3) + p_2(2) + p_2(1) + p_2(0) \\ &= 4 + 2 + 2 + 1 + 1. \end{aligned}$$

We ran our tree isomorphism program on some other popular graphs, namely Q_3 and the Petersen graph. Let Q_n denote the n -dimensional cube and let P denote the Petersen graph. $\tau(Q_n)$ is known, the values $\tau(Q_3) = 384$ and $\tau(P) = 2000$ are generally known, however, it appears that $I(Q_n)$ and $I(P)$ may not be so universally known. After applying our algorithm to Q_3 and P , we have found that $I(Q_3) = 6$ and $I(P) = 20$. Table 2 gives the breakdown of the size of each isomorphism class in Q_3 . For example, row 2 denotes that there are 3 classes, each containing 48 trees. Table 3 gives drawings of a representative tree from each of the 6-classes. We remark the class containing the spanning paths has 72 trees. Table 4 gives the different class sizes for the Petersen graph. On Austin Mohr's website [7], there are drawings of representative trees for the Petersen graph similar to Table 3. There are also drawings for the trees given in Table 1.

$$T = 384$$

$$I = 6$$




Distribution of Class Sizes

	Num Trees	Size of Class
	24	1
	48	3
	72	1
	144	1
Total	384	

Table 2: Q_3

$$T = 384$$

$$I = 6$$




48 in class	48 in class
144 in class	48 in class
72 in class	24 in class

Table 3: Q_3

$$T = 2000$$

$$I = 20$$



Distribution of Class Sizes

	Num Trees	Size of Class
	10	1
	30	1
	40	1
	60	4
	120	12
	240	1
Total	2000	20

Table 4: Petersen Graph

References

- [1] N.H. Abel, Beweis eines Ausdrucks, von welchem die Binomial-Formel ein einzelner Fall ist. *J. reine angew. Math.* **1**, (1826), 159–160. Reprinted in *Œuvres Complètes, 2nd ed.*, **1**, (1881), 102–103.
- [2] T.L. Austin, The enumeration of point labelled chromatic graphs and trees, *Canad. J. Math.* **12**, (1960), 535–545.
- [3] L. Clark, On the enumeration of multipartite spanning trees of the complete graph, *Bull. of the ICA* **38** (2003), 50–60.
- [4] O. Eğecioğlu and J.B. Remmel, A bijection for spanning trees of complete multipartite graphs, *Congressus Numerantium* **100** (1994), 225–243.
- [5] I.J. Good, The generalization of Lagrange’s expansion and the enumeration of trees, *Proc. of the Cambridge Phil. Soc.* **61** (1965), 499–517.
- [6] R.P. Lewis, The number of spanning trees of a complete multipartite graph, *Discrete Math.* **197/198** (1999), 537–541.
- [7] A. Mohr, <http://www.cs.siu.edu/~amohr/>.
- [8] J.W. Moon, Counting labeled trees, *Canad. Math. Monograph No. 1*, Canad. Math. Congress, 1970.
- [9] R. Otter, The number of trees, *Ann. of Math.* **49** (1948), 583–599.
- [10] T.D. Porter, Generating the list of spanning trees in $K_{s,t}$, *J. of Comb. Math. and Comb. Comput.* **50** (2004), 17–32.
- [11] L. Székely, Abel’s binomial Theorem, <http://www.math.sc.edu/~szekely/abel.pdf>.
- [12] D.B. West, *Introduction to Graph Theory*, Second Edition, Prentice-Hall, Upper Saddle River, NJ, 2001.